

ARE YOU
ADDISON'S
AWARE?

ZYCORTAL®

HIDDEN DISEASE. VISIBLE ANSWER.

Using Zycortal in your practice

No previous treatment

Fludrocortisone alone

Fludrocortisone and Prednisolone

Day 1 for all cases - Administer starting dose of 2.2 mg/kg subcutaneously

Day 1 - Administer 0.2-0.4 mg/kg SID of Prednisolone alongside the Zycortal

Day 1 - Administer Fludrocortisone tablets as normal

Day 1 - Administer Fludrocortisone and Prednisolone tablets as normal

Day 2 - Halve the original dose of Fludrocortisone tablets and administer 0.2-0.4 mg/kg SID of Prednisolone

Day 2 - Halve the original dose of Fludrocortisone tablets. Increase Prednisolone to 0.2-0.4 mg/kg SID (if not already at this dose)

Day 3 - Quarter the original dose of Fludrocortisone tablets. Continue with Prednisolone

Day 3 - Quarter the original dose of Fludrocortisone tablets. Continue with Prednisolone

Day 4 - Stop Fludrocortisone. Continue with Prednisolone

Day 4 - Stop Fludrocortisone. Continue with Prednisolone.

All dogs should then receive a clinical exam and blood electrolyte check at day 10 and day 28.

ZYCORTAL: Zycortal contains desoxycortone pivalate

Dechra Veterinary Products Limited, Sansaw Business Park, Hadnall, Shrewsbury, Shropshire SY4 4AS. Tel: +44 (0)1939 211200 Fax: +44 (0)1939 211201 www.dechra.co.uk

Registered Office: 24 Cheshire Avenue, Cheshire Business Park, Lostock Gralam, Northwich CW9 7UA.

Registered in England and Wales, Company Registration No.5385888. Dechra Veterinary Products Limited is a trading business of Dechra Pharmaceuticals PLC.

Please refer to package inserts UK: POM-V IE: POM

Use medicines responsibly: www.noah.co.uk/responsible

www.dechra.com/xxxx ©Dechra Veterinary Products A/S March 2023


Dechra