

## Suitable dogs

- Once- or twice-daily Vetoryl dosing
- Adrenal- or pituitary-dependent hyperadrenocorticism (HAC)
- Clinically well dogs (with or without signs of HAC)
- Calm dogs

## Unsuitable dogs

- Aggressive dogs
- Stressed dogs (e.g. persistently barking)
- Unwell dogs



**Better teamwork. Better results.**

The more we work with owners  
the better the outcome for their dog.

## Appointment

- Book appointment just before the next Vetoryl dose is due
- If the dog is normally given Vetoryl at an inconvenient time then ask owner to give at a convenient time from at least the day before
- Make sure owner has not given Vetoryl and that nothing stressful has happened that morning (e.g. vomiting, injury)
- When re-evaluating a patient receiving Vetoryl, it is important to carefully consult with the owner regarding their dog's clinical response at home

## Sample

- Take sample immediately after examination and before administration of Vetoryl
- 1 to 2 ml of blood in heparin or serum tube
- Can be separated and stored for up to 1 week
- Send to an external laboratory participating in an external quality assurance scheme (e.g. ESVE- or SCE programmes) and preferably that uses a Siemens IMMULITE® – or a method that has been validated against this machine

<sup>1</sup> Polyuria/polydipsia, polyphagia, panting & lethargy should resolve or improve in 1 month. Alopecia & pot belly appearance may take 3-6 months to resolve

<sup>2</sup> Reassess history and consider Pre-Vetoryl ACTH Stimulation Test. Contact Dechra Technical Services for further information

<sup>3</sup> If symptoms are not adequately controlled for an entire 24 hour inter-dose period, consider increasing the total daily dose by up to 50% and dividing it equally between morning and evening doses

<sup>4</sup> Use combinations of capsule sizes to increase or decrease the once or twice daily dose

<sup>5</sup> Dexamethasone to treat hypocortisolaemia, IV 0.9% NaCl to resolve dehydration & hyperkalaemia; alternatively hydrocortisone CRI and IV 0.9% NaCl

Since the Pre-Vetoryl Cortisol test does not feature on the Vetoryl Summary of Product Characteristics, informed owner consent for 'off-label' monitoring should be obtained.

**VETORYL:** Vetoryl contains trilostane UK: **POM-V** IE: **POM**

For further information contact: Dechra Veterinary Products Limited, Sansaw Business Park, Hadnall, Shrewsbury, Shropshire SY4 4AS T +44 (0)1939 211200 F +44 (0)1939 211201 [www.dechra.co.uk](http://www.dechra.co.uk) [www.dechra.ie](http://www.dechra.ie)  
Registered Office: 24 Cheshire Avenue, Cheshire Business Park, Lostock Gralam, Northwich CW9 7UA.  
Registered in England and Wales, Company Registration No.5385888. Dechra Veterinary Products Limited is a trading business of Dechra Pharmaceuticals PLC


Use medicines responsibly: [www.noah.co.uk/responsible](http://www.noah.co.uk/responsible)

Dec 2023 DVP1412web



# How to monitor Cushing's

Developed by Ian Ramsey BVSc, PhD, DSAM, Dipl. ECVIM-CA, FHEA, MRCVS, Federico Fracassi DVM, PhD, Dipl. ECVIM-CA, Nadja Sieber-Ruckstuhl PhD, Dr. med. vet, Dipl. ACVIM, Dipl. ECVIM-CA



\*These figures are based on the use of Siemens IMMULITE 1000/2000 analysers: other analysers may have different thresholds and veterinary surgeons should consult their laboratory if in doubt.